

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ PREVATHON™ INSECT CONTROL

EPA Reg. No. 352-844

FOR CONTROL OF GRASSHOPPER NYMPHS AND SUPPRESSION OF GRASSHOPPER ADULTS IN GRASS FORAGE FODDER AND HAY (RANGELAND & PASTURE GRASS) IN THE STATE OF TEXAS

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DuPont™ PREVATHON™ insecticide is recommended for the control of grasshopper nymphal stages and suppression of grasshopper adults in grass forage, fodder, and hay: any grass, Gramineae family (rangeland & pasture grass).

Apply PREVATHON™ at 8.0 to 16.0 fluid ounces per acre as a foliar spray via aerial application or ground application. Apply 16.0 fluid ounces per acre for suppression of adult grasshoppers. For best results with foliar sprays, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v).

Apply when grasshopper nymph populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results applications should be made when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs.

PREVATHON™ may be applied using a RAAT application technique. Read and follow all spray drift management guidance in the Federal Section 3 PREVATHON™ label.

Once grasshopper nymphs contact and/or ingest PREVATHON™ there will be rapid feeding cessation; full expression of mortality may take a week or more.

Do not make more than two sequential applications of PREVATHON™ before rotating to another registered insecticide having a different mode-of-action.

PRECAUTIONS

Do not apply more than 60.0 fluid ounces of PREVATHON™ per acre per crop.

Do not make more than 2 applications per acre per crop. The minimum retreatment interval is 7 days.

The preharvest interval is 0 days. The field re-entry interval is 4 hours.

Read and follow all applicable directions, restrictions, and precautions on the EPA-registered label.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2011 E.I. du Pont de Nemours and Company, 1007 Market Street, Wilmington, DE 19898